



Understanding Your Diabetes

- The Trust offers support, understanding and information to people with diabetes and those who care for them.
- We listen to the needs of people who live with diabetes and do our utmost to offer help and support.
- We raise awareness of important issues for people living with diabetes and lobby governments on issues that affect people's lives.
- We fund research into ways of improving the lives of people with diabetes

Facts about Diabetes

Diabetes is a chronic disease that affects over 3.5 million people in the UK, around 90% of whom have Type 2 diabetes. In the UK there are 23,000 children with Type 1 diabetes under the age of 15 years.

There are several forms of diabetes but the main ones are Type 1 and Type 2 diabetes

Type 1 diabetes

- This type accounts for about 10 to 15% of the total number of people with diabetes. It usually affects children and adults up to the age of 40 but can be diagnosed in much older people. The number of children diagnosed under the age of 5 is markedly increasing. It is also referred to as insulin dependent diabetes or juvenile diabetes.
- Type 1 diabetes is caused by the body's immune system attacking the insulin producing beta cells in the pancreas. The body no longer produces insulin and glucose levels rise. Treatment with insulin injections is always required for survival. It is usually diagnosed as an acute condition.
- It is estimated that between 10,000 and 20,000 people are treated with natural animal insulin and the remainder with synthetic 'human' or analogue insulins made by genetic engineering.
- There is no cure for Type 1 diabetes and cause has not been established. It is thought that there may be several causes with a genetic link in some people. Research shows that a common virus may trigger the body's immune system to attack its own insulin-producing pancreatic cells.

Type 2 diabetes

- This type of diabetes affects about 90% of the total number of people with diabetes, nearly 3.5 million people and it is thought that there are around 800,000 people undiagnosed. It occurs mainly in people over the age of 40. However, with increasingly levels of obesity, it is becoming increasingly common in younger people. Type 2 diabetes is also referred to as non-insulin dependent diabetes.
- In Type 2 diabetes, the pancreas often still produces some insulin but either not enough or it is not used properly by the various organs in the body [so there can be too much insulin in the system].
- Type 2 diabetes can sometimes be treated with diet and exercise alone, sometimes with oral blood glucose lowering drugs and if this fails to reduce blood glucose levels sufficiently, then treatment with injectable drugs or insulin is necessary. On average people with Type 2 diabetes start to take insulin 7 years after diagnosis.
- Type 2 diabetes can remain undiagnosed for several years during which time the blood glucose levels are too high causing damage and some of the complications of diabetes. People with Type 2 diabetes are often diagnosed as a result of having complications rather than because they suspect they have Type 2 diabetes.
- There is a tendency for Type 2 diabetes to run in families but a sedentary lifestyle and being overweight or obese are also causes.



Gestational diabetes

This type of diabetes may occur during pregnancy but disappears after the baby is born. Like other forms of diabetes, gestational diabetes affects the way the body uses glucose.

If gestational diabetes is untreated or uncontrolled, it can result in a variety of health problems for both the mother and baby. So it is important that a treatment plan is worked out to keep blood sugars within the normal range. The good news is that controlling blood sugars can help to ensure a healthy pregnancy and a healthy baby.

Latent Autoimmune Diabetes [LADA]

Sometimes people don't really know whether they have Type 1 or Type 2 diabetes and no one has actually told them! It could be that they have latent autoimmune diabetes [LADA] which is also called Late-onset Autoimmune Diabetes of Adulthood, Slow onset Type 1 diabetes or Type 1.5 diabetes.

People with LADA are usually diagnosed at a later age than people with typical Type 1 diabetes. It is now thought that 20% of people diagnosed with Type 2 diabetes could actually have LADA.

Unlike Type 2 diabetes, people with LADA do not have insulin resistance and are not obese or overweight at diagnosis. The characteristics of LADA are:

- **Adult age at diagnosis – usually over 25 years.**
- **People are often mistakenly thought to have non-obese Type 2 diabetes because of their age at diagnosis and the fact they initially respond to treatment with diet with or without tablets.**

- Treatment with insulin gradually becomes necessary and often within months of diagnosis. Some specialists treat LADA with insulin immediately but it is not known whether this early treatment is beneficial for the remaining insulin-producing beta cells.
- They have low C-peptide levels.
- They are unlikely to have a family history of Type 2 diabetes.

Secondary diabetes

This is where diabetes is caused by some outside source. Some drugs can cause diabetes, such as steroids and some antidepressants or it can be caused by surgery, such as removal or partial removal of the pancreas.

Pre-diabetes

- Pre-diabetes is a relatively new term for when blood glucose levels are higher than normal but not sufficiently high to be classed as diabetes.
- Most people at risk of Type 2 diabetes do not have any symptoms but if they are overweight and over 45 years old, it may be recommended that they are tested for diabetes.
- Type 2 diabetes can be prevented, delayed or even reversed by weight loss of 5 to 10% of the starting weight and by being physically active.

Rare forms of diabetes

There are rare forms of diabetes such as MODY and Diabetes Insipidus.

Whichever type of diabetes you have, the more you know about your condition and how to manage it, the better you will feel and the better your health will be.

The term 'diabetes' blurs the lines and causes confusion

As we can see, Type 1 and Type 2 diabetes are two very different conditions - they have different causes and affect different groups of people. All too often they are put together under the general umbrella of 'diabetes' - in press articles and even NHS information and diabetes organisations' magazines are guilty of just referring to 'diabetes'. This blurring of the lines between Type 1 and Type 2 can cause unnecessary fears, confusion and misunderstandings that can be dangerous. The misconception that 'diabetes' is simply a lifestyle condition minimises the public perception of the seriousness of both types of diabetes.

Why blood glucose levels need to be controlled

If blood glucose levels are too high then this can lead to long-term complications. Type 1 and Type 2 diabetes are different diseases in cause, effect and treatment but the same long-term complications can arise in both types of the condition. The risk of complications is reduced by the treatment of diabetes and by lifestyle changes to reduce blood glucose levels. In the case of Type 2 diabetes, early diagnosis helps to reduce the risk of complications developing.

The recommendations are that people with both types of diabetes should try to keep their blood glucose levels as near normal as possible, that is between 4 and 7mmols/l. However, it is well recognised that in Type 1 diabetes this increases the risk of severe hypoglycaemia threefold so good control is a balance between achieving blood sugars that are as near normal as possible but at the same time avoiding hypoglycaemia. Hypoglycaemia in people treated with insulin is caused by the blood glucose lowering effect insulin itself and not by diabetes. For many people, hypoglycaemia and the avoidance of it, is one of their major daily concerns.

The complications affect:**The eyes**

Diabetes can affect the blood vessels at the back of the eye [retinopathy] and this can lead to visual impairment or blindness.

The heart and vascular system


Diabetes can affect the heart and the vascular system making people more susceptible to heart disease and stroke. It can also cause blood clots in the vessels in the legs which may result in amputation. Amputations are 50-80 times higher in people with diabetes than the general population.

Kidneys

Diabetes can affect the kidneys resulting in damage or kidney failure [nephropathy].

Nerves

Diabetes may cause nerve damage [neuropathy]. The most common form of nerve damage is in the extremities leading to pain or loss of sensation in the feet and ulceration of the legs. Again, this can lead to amputation.



Understanding diabetes and its treatment

To understand diabetes and its treatment, we first have to learn how the body works in people without diabetes.

Normal metabolism

Normally during food digestion the body breaks down the carbohydrates you eat into simple sugars, known as glucose. The glucose is absorbed into the blood and transported around the body by the blood vessel system to provide the energy needed for all our activities. As food is eaten, insulin is released into the blood stream and this allows glucose to enter the body's muscle, fat and liver cells.

The pancreas

The pancreas is a gland behind the stomach. Beta cells in the pancreas produce the hormone insulin to help to control the levels of glucose in the blood.

The liver

The liver also plays a part in maintaining normal blood glucose levels. When there is more glucose in the cells than your body needs for energy, it is removed from the blood and stored in the liver as glycogen. It can then be used when necessary, such as at times when you run low on glucose eg if you have missed a meal. In such situations the liver releases glucose into the bloodstream.

What affects the amount of glucose in the blood?

The amount of glucose in the blood varies according to several factors – the food eaten, exercise, stress and infections. The relationship between insulin production, glucose and the liver makes sure that the blood glucose levels stay within normal limits. For people without diabetes, these are 4 to 7mmols/l.

What happens in Type 1 diabetes...

In people with Type 1 diabetes, the beta cells in the pancreas cannot produce insulin. When carbohydrates are eaten, no insulin is produced and so the glucose levels in the blood rise higher and higher.


The body cannot cope with this and so the excess glucose is passed through the body into the urine. This means that people with untreated diabetes pass urine frequently to get rid of the excess glucose. In turn, this makes them thirsty because the body gets dehydrated. *These are the classic signs of undiagnosed diabetes – thirst and peeing a lot.* The body becomes short of energy as a result of the glucose being excreted and the person feels tired. The body starts to burn its own fats to provide the necessary energy and there is weight loss. Treatment is essential at this stage and it is often an acute emergency situation. Type 1 diabetes always requires treatment with insulin.

What happen in Type 2 diabetes...

In Type 2 diabetes either the insulin-producing cells in the pancreas do not produce enough insulin or the insulin they produce is not used properly – this is called insulin resistance. The blood glucose levels gradually rise and this may be so gradual that it takes place over several years before diagnosis.

As the onset is gradual, the extreme symptoms that occur with Type 1 diabetes don't occur but they creep on gradually. Blood glucose levels may not rise sufficiently high for glucose to spill into the urine, so Type 2 diabetes can go undiagnosed during which time, it can be causing some of the complications.

Three important factors in the treatment of Type 1 and Type 2 diabetes



Although Type 1 and Type 2 diabetes are very different conditions, the treatment of both types of diabetes has three factors in common – medication [insulin or tablets], diet and exercise. All three are important, they all affect blood sugar levels and they all affect each other. So in both Type 1 and Type 2 diabetes there is an important relationship between medication [insulin or tablets], diet and exercise and to achieve good diabetic control we have to try to balance all three. It is much easier to manage both types of diabetes if we understand about carbohydrates, about exercise and about insulin or tablets.

Understanding Carbohydrates


Carbohydrates are sugars and starches – bread, potatoes, rice, pasta, cereals and sugar. They provide the energy our bodies need for all its various activities. [Energy is also supplied by fats.]

How does the body convert carbohydrates we eat into glucose?

- When we eat, beta cells in the pancreas produce insulin, a hormone which controls the glucose levels in the blood.
- The pancreas produces the right amount of insulin for the amount of carbohydrates we have eaten and this keeps the level of glucose in the blood within the normal range.
- If we have eaten more carbohydrates than we need for energy at a particular time, then the excess glucose is stored in the liver. This is used for emergencies such as extra physical activity, or occasions when the blood glucose levels drop unexpectedly, such as times of fear.

The type of carbohydrate we eat is important to people with diabetes:

- Sugary foods, such as cakes and biscuits, will raise blood sugars more quickly and higher in people with diabetes and more injected insulin or medications may be necessary. Sugary carbohydrates tend not to last as long in the body so blood sugars may drop before the next meal. Sugary carbohydrates tend to make blood glucose levels peak and trough.
 - Slow-acting carbohydrate, those such as cereals and bread, do not raise the blood sugars as quickly or as high after eating. They last longer and therefore tend to give more even blood glucose levels. The amount of insulin, or medication (in Type 2 diabetes) that is needed, may be less.



Understanding exercise

Physical activity and diabetes - it is recommended that we all take exercise – at least 30 minutes brisk walking, 5 days a week.

Facts about exercise

- People with diabetes are at greater risk of coronary heart disease and regular exercise reduces this risk.
- Physical activity can reduce the amount of medication needed or reduce the daily insulin intake.
- Moderate, rhythmic exercise seems to reduce the risk of people developing Type 2 diabetes in middle age.
- Physical activity appears to raise HDL (good) cholesterol levels but it does not affect LDL (bad) cholesterol levels.
- It lowers blood pressure or prevents it from developing.
- It helps to reach and maintain a healthy weight.

Types of activity

Aerobic activity – this type of exercise benefits your heart. It is any activity that is rhythmic and repetitive eg walking, swimming, cycling or dancing, which increase the body's demand for oxygen so making the heart and lungs work harder and more efficiently.

Isometric exercise - this increases muscle tension without moving a joint eg pushing against a wall. Isometric exercise does not help the heart and



circulation. It should be avoided by people with heart disease or high blood pressure because it can increase blood pressure and put the heart under stress.

Exercise and the risk of hypoglycaemia (low blood glucose levels)

Keeping fit is recommended for people with diabetes, including those being treated with insulin. However, injected insulin cannot mimic the response of a healthy pancreas to exercise.

If you have diabetes which is treated with insulin, or with one of the Type 2 diabetes drugs that can cause hypoglycaemia, then exercise can cause hypoglycaemia. Hypoglycaemia can also occur 12 – 14 hours or even longer after exercise. This is because the body uses up any circulating glucose to try to replace the glycogen stores in the liver that were used up during exercise. In addition, exercise increases the sensitivity of the body tissues to insulin, especially the muscles.

It is important to eat sufficient carbohydrates before, during and after exercise to avoid hypoglycaemia by:

- Eating a meal of slow-acting carbohydrates about an hour before exercising will keep your blood sugars steady during exercise. Examples: porridge, cereal or multi-grain bread.
- Eating fast-acting carbohydrate immediately after exercise will help to prevent hypoglycaemia and will help to re-stock the liver stores of glycogen which the body turns into glucose when needed. Examples: a piece of fruit, fruit juice or biscuits.
- Regular blood glucose monitoring is important when exercising to avoid both high and low blood sugars.

Understanding carbohydrates and exercise is necessary in both types of diabetes but the medications for Type 1 and Type 2 diabetes are different so we have to look at Type 1 and Type 2 diabetes separately.

Type 1 diabetes – insulins and insulin regimes

Types of insulin (also known as species of insulin)

- For nearly 60 years all people with diabetes who required insulin treatment used animal insulin. Originally this was beef insulin but in the 1970s highly purified pork insulin also became available. All insulins are now highly purified whether beef, pork or genetically modified human and analogue insulin (GM insulin).
- In 1982 genetically modified so-called 'human' insulin was introduced and about 85% of people using insulin were transferred to human insulin on the assumption, not proof, that it was better than animal insulin.
- In 1998 the first insulin analogue was introduced. Analogue insulins are made from 'human' insulin by genetically modifying it again.

So in the UK, there is a choice of analogue, animal and human insulins and all are available on an NHS prescription.

Action and duration times of insulins

In Type 1 diabetes usually between 2 - 4 daily injections are given but even so, this does not mimic the body's normal action of insulin being produced according to need [that is the amount of carbohydrate eaten]. Therefore, the amount insulin, its peak of activity and the duration of its action have to be balanced with the amount of food eaten and the level of activity or energy required.

- Rapid and short-acting insulins deal with the carbohydrates eaten at meal times.
- Intermediate and long-acting insulins work in the background all the time.

Table 1 shows the different insulins according to their origin and their length of action.

INSULIN	RAPID	SHORT	INTERMEDIATE	LONG
Analogue	Apidra Humalog NovoRapid			Abasaglar Lantus Levemir Toujeo (300units/mL) Tresiba
Animal		Hypurin Bovine Neutral Hypurin Porcine Neutral	Hypurin Bovine Isophane, Hypurin Porcine Isophane	Hypurin Bovine lente Hypurin Bovine PZI
Human		Actrapid, Humulin S, Insuman Rapid	Humulin I, Insuman basal, Insulatard	

Also available are pre-mixed insulins of rapid-acting or short-acting insulins with an intermediate-acting insulin. These are usually used where only two injections are day are required.

Table 2 shows the different pre-mixed insulins.

INSULIN	Pre-mixed
ANALOGUE	Humalog Mix 25 and Mix 50, NovoMix 30
ANIMAL	Hypurin Porcine 30/70
HUMAN	Humulin M3 and Insuman Comb 15, 25 and 50

Table 3 is a rough guide to the activity curves of the different insulins. It is important to remember that these times of peak activity and duration are general and will vary in different people. This table is listed in order of the development of the different insulin types.

Table 3

INSULIN TYPE	ONSET	PEAK [hours]	EFFECTIVE DURATION [hours]	MAXIMUM DURATION [hours]
ANIMAL				
Regular [short]	0.5 - 2 hours	3 - 4	4 - 6	6 - 8
NPH [intermediate]	4 - 6 hours	8 - 14	16 - 20	20- 24

'HUMAN'				
Regular [short]	0.5 - 1 hour	2 - 3	3 - 6	6 - 10
NPH [Intermediate]	2- 4 hours	4 - 10	10 - 16	14 - 18
ANALOGUES				
Rapid	Immediate	Immediate to about 1 hour	Tails off from peak over 3- 4 hours	About 4 hours
Abasaglar (long)	2 hours	None	24hours	
Lantus [long]	2 hours	None	24hours	
Levemir [long]	2 hours	None	14 or more hours	
Toujeo (long)			Once daily at the same time	
Tresiba (ultra-long)			Once daily, lasts beyond 42 hours	

Note: Long-acting beef insulins [Bovine lente and Bovine PZI] are available but they are not included in this table as they are rarely used.

Key facts about insulins and their actions:

- The manufacturers' profiles of the speed of onset, peak of action and duration of action are only a rough guide and may vary. Insulins act differently in different people.
- The speed of action of injected insulin varies at different injection sites. For example, if Hypurin Porcine Neutral is injected above the umbilicus [tummy button] its speed of action is much quicker, and similar to a rapid-acting analogue, than if it is injected below the tummy button.
- The depth of the injection may also affect the speed of action and it is important to have the correct sized needle.



Insulin Regimes

Standard insulin regimes include:

Short or rapid-acting insulin at meal times with intermediate or long-acting insulin.

- Rapid-acting insulin starts to act almost immediately and was designed to lower post-meal blood sugars but short-acting insulin takes between 20 to 30 minutes to start acting. However, short-acting insulin has the advantage of lasting longer than the rapid-acting insulins and it is still active by the next meal time. It is worth noting that for people who eat a high fibre diet of slow-acting carbohydrate, rapid-acting insulin may be too rapid acting as it will work before the carbohydrate has got into the system and this could result in hypoglycaemia.
- Intermediate insulins provide the background [basal] insulin and last for around 12 hours and therefore are best given twice a day [before breakfast and the evening meal] to give 24 hour background insulin cover.
- The long-acting analogues, Lantus, Levemir and Abasaglar, differ in the duration of action. According to the manufacturers, the duration of action of Lantus and Abasaglar is 24 hours and the duration of Levemir is 12 – 14 hours (the manufacturers advise that it is a once or twice daily insulin.) Toujeo is a long-acting insulin but has a higher strength of 300units/L. Tresiba is an ultra-long-acting insulin which lasts beyond 24 hours.

Pre-mixed insulins morning and evening.

These insulins are usually used twice a day. As they are a combination of 30% short-acting or rapid-acting and 70% intermediate-acting insulin given twice daily, they provide 24 hour basal insulin coverage. The drawback to pre-mixed insulins is less flexibility because the short/rapid-acting insulin cannot be adjusted independently of the longer-acting insulin.

Long-acting insulins alone

These are sometimes used but usually for Type 2 diabetes either once or twice daily depending on the type of insulin being used.

Note: Variations on these standard regimes can be introduced to suit individual lifestyles or to try to obtain more even blood glucose levels where people feel able to manage more complex regimes.


Here are some examples:

- Pre-mixed insulins can be used with the addition of long or intermediate-acting insulins.
- If rapid-acting analogue insulin is too short-acting so that blood sugars go up before the next meal, a short-acting insulin with a longer duration can be given at the same time.
- If Hypurin Porcine Neutral is the short-acting insulin normally used but this does not act quickly enough after meals, a small dose of a rapid-acting insulin analogue can also be given with the meal. This may only apply with certain mealtime injections eg before breakfast if blood sugars are high in the morning.
- Small doses of short-acting insulin [human or animal] or rapid-acting analogue insulin can be given at any time of day to bring down unexpected high blood sugars. Alternatively taking exercise [brisk walk or cycle ride] can reduce the high blood sugar which may be preferable to giving extra insulin.

Insulin pump therapy

Some children and adults with Type 1 diabetes use insulin pump therapy [continuous subcutaneous insulin infusion or CSII] instead of injections.

This is where there is a continuous infusion of short/rapid-acting insulin delivered by a short cannula under the skin with boluses of insulin given to cover meals.



Your choices of insulin and insulin regimes

NHS policy is to provide patient-centred care, defined by National Institute for Health and Clinical Excellence [NICE] as treatment that takes into account patients' need and preferences. NICE goes on to say that 'good communication' is essential, supported by evidence-based information, to allow patients to reach informed decisions about their care. [NICE Guidelines for Type 2 diabetes]

Type 1 diabetes is a condition that is largely self-managed, so it is especially important that you are involved and can make an informed choice of insulin and regime to suit your needs, your lifestyle and your preferences. Type 1 diabetes is also a condition that affects the people around you, your family or work colleagues and so it is important that they have the opportunity to be involved in your choices, if that is what you wish.

Factors that may influence your choices

There is a choice of insulins - animal, human or analogue insulins.

Diet: The type of diet eaten may affect your choice of insulin. Here are just some examples:

- You may prefer a low carbohydrate diet to reduce your daily intake of insulin and reduce the risks of severe hypos by only needing small doses of insulin.
- If you prefer a low carbohydrate diet it may mean that a lunchtime injection is not necessary eg if you eat salad with no carbohydrate.
- If meals are largely slow-acting carbohydrates, then animal or human insulin may be better than a rapid-acting analogue which acts too quickly and does not last long enough for the slow-acting carbohydrate.

Practicalities, lifestyle and quality of life

Your lifestyle may be such that it influences your choice of insulin regime. Here are just some examples:

- If your job involves a lot of driving, you may consider that it is safer for you to run your blood sugars less tight, than the target of near normal, so that you do not increase your risks of being hypo while driving.
- If you live alone, you may prefer to run your night time blood sugars a little higher to avoid the risk of a night hypo.
- If injecting at lunchtime is a problem, such as for children at school, then you may choose to use short-acting insulin and twice daily intermediate insulins that cover lunchtime carbohydrate eaten without the need for an injection.
- If you find multi-dose regimes difficult and confusing, you may find it easier to use a twice daily injection regime to give you a better quality of life.

Adverse reactions, safety and efficacy

As with all new drugs, when new insulins are introduced to the market, it is important to be aware that they have only been trialed in a relatively small number of people, so they may not suit everyone. Adverse reactions may occur in some people but may not show up until the new insulin is used in the wider diabetic population. In addition, the long-term safety and efficacy of new insulins is unknown, so you may prefer to stay with a previous insulin that has a history of safety and where any adverse reactions are known.

You have a choice of treatment and so IDDT recommends that you discuss your choice and the risks and benefits with your doctor or diabetes team.

The International Diabetes Federation Position Statement, March 2005 [accessed April 2012] states:

“There is no overwhelming evidence to prefer one species of insulin over another and patients should not be changed from one species to another without reason.”

Type 2 diabetes treatment

As we have established, essential parts of the treatment of Type 2 diabetes are diet and exercise and appropriate medications, when necessary, to try to achieve target blood glucose levels. NICE guidelines recommend that target blood glucose levels for Type 2 diabetes should be agreed with you, the patients but unlike Type 1 diabetes, NICE also recommends that pursuing highly intensive management to levels of less than 48mmols/mol (6.5%) should be avoided.

Diet and exercise alone

Initially for some people, diet alone with exercise may produce the desired lowering of blood glucose control. However this may not be appropriate for everyone and may not maintain blood glucose control so drugs have to be used.

Type 2 drugs

There are a range of drugs to treat Type 2 diabetes all of which should be used with diet and exercise. There are three main drugs that are used to treat Type 2 diabetes when diet and exercise alone are not sufficient. NICE recommends that they are used as follows:

NICE recommends that they are used as follows:

- **Biguanides (Metformin)** – this reduces insulin resistance, which means that it improves the body's ability to use the insulin that is still being produced. It does not increase insulin production and therefore does not cause hypoglycaemia [low blood sugars]. It is particularly used in people who are overweight. Its main adverse effects are stomach upsets. If metformin alone does not control blood glucose levels then another drug may be added from the range known as sulfonylureas.
- **Sulfonylureas (Glibeclamide, Gliclazide, Glimepiride, Glipizide, Tolbutamide)** – these drugs stimulate the production of insulin and therefore may cause hypoglycaemia.

- **Glitazones (Pioglitazone)** can be added to treatment. This drug works by reducing insulin resistance in fat tissue, muscles and the liver. Research has shown that this family of drugs can cause fluid retention, heart and liver problems and increase the risk of osteoporosis and some doctors are no longer prescribing these drugs. It is advisable to discuss them with your doctor.

There are other drugs available to treat Type 2 diabetes which your doctor may discuss with you. There are also injectable drugs, exenatide (Byetta/Bydureon) and liraglutide (Victoza). Although these are injected, they should not be confused with insulin. The most common side effects are stomach upsets and the main advantage of these drugs is that they can help people to lose weight.

You should remember that if you are taking any medication that can cause hypos then you should check your blood sugar levels regularly, particularly if you drive. You should discuss this with your doctor.

Addition of insulin

If the above tablets fail to control blood sugars sufficiently, then insulin may be required. NICE recommendations for insulin regimes in this situation are as follows:

- **Begin with intermediate-acting insulin at bedtime or twice daily according to need.**
- **Alternatively, there are situations where a once-daily long-acting insulin analogue may be advisable: [i] if assistance with injections is needed, [ii] if there is significant and frequent hypoglycaemia or [iii] twice daily basal insulin plus oral medications would otherwise be needed.**
- **Consider twice daily pre-mixed human insulin especially where the HbA1cs are greater than 75 mmol/mol] (9%). A once daily regime with pre-mixed insulin may be an option.**

- Consider pre-mixed insulin analogues rather than human pre-mixed analogues when: [i] immediate injection before a meal is preferred, [ii] hypoglycaemia is a problem or [iii] there are marked rises in blood glucose levels after a meal.

These are basic insulin regimes that your doctor may recommend for you to discuss. However, structured education should also be offered to you and/or your family carer at the time of diagnosis to help you to learn to manage your diabetes.

NICE recommends this education programme should include:

- Continuing telephone support and support from an appropriately trained and experienced healthcare professional.
- Frequent self-monitoring.
- Dose titration to target [how to adjust insulin doses].
- Dietary understanding.
- Management of hypoglycaemia – how to avoid it and treat it.
- Management of acute changes in glucose control.

Summary of the key NICE Guidance for Type 2 diabetes for implementation by health professionals

Patient education

- Offer structured education to every person and/or their carer at and around the time of diagnosis – this can be in group sessions run by people who have been specially trained to do this.
- Provide individualised and ongoing nutritional advice from a healthcare professional with specific expertise and competencies in nutrition.

Setting a target HbA1c

- Involve the person in decisions about their individual HbA1c target level, which may be above that of 48mmol/mol (6.5%) set for people with Type 2 diabetes in general.
- Encourage the person to maintain their individual target unless the resulting side effects [including hypoglycaemia] or their efforts to achieve this impair their quality of life.
- Offer therapy [lifestyle and medication] to help achieve and maintain the HbA1c target level.
- Inform a person with higher HbA1c that any reduction in HbA1c towards the agreed target is advantageous to future health.
- Avoid pursuing highly intensive management to levels of less than 48mmol/mol (6.5%)

Self-monitoring

- Offer self-monitoring of plasma glucose to a person newly diagnosed with Type 2 diabetes only as an integral part of his or her self-management education. Discuss its purpose and agree how it should be interpreted and acted upon.

Regular Reviews

Looking after diabetes, whether you have Type 1 or Type 2, means having regular reviews with your doctor and/or diabetes team to help you control your blood glucose levels and also to check for any diabetes complications. It may be that you will be prescribed other drugs to help you, such as for blood pressure or raised cholesterol levels. This is also your opportunity to discuss any concerns or questions you may have and this way you are working together to provide the best care of you and your diabetes.



HbA1c Converter

The HbA1c test measures your average blood glucose levels over the last 6 to 8 weeks. The results of the HbA1c used to be a percentage figure but the measurements have changed in the UK to mmol/mol. Some people and some articles are still using the old measurements, so it is important that we are aware of both.

Here is a table showing the DCCT measurements as a percentage, the new IFCC measurements, and the average blood glucose measurements you can expect to be associated with each particular level of HbA1c

HbA1c (DCCT) measurement (%)	HbA1c (IFCC) Measurement from June 2011 (mmol/mol)	Average blood glucose level for this HbA1c, mmol/l
6	42	7.0 (range 5.5-8.5)
7	53	8.6 (range 6.8-10.3)
8	64	10.2 (range 8.1-12.1)
9	75	11.8 (range 9.4-13.9)
10	86	13.4 (range 10.7-15.7)
11	97	14.9 (range 12.0-17.5)
12	108	16.5 (range 13.3-19.3)
13	119	18.6 (range 14.6-21.1)

The 9 Key Tests

This chart is to remind you of the 9 tests which should be carried out at your annual review. You can keep record of which tests were carried out by completing the date and ticking the box for each test. At your review, ask if all 9 tests have been done and if not, ask why not. If necessary, ask for the tests to be carried out.

Date	Weight	Blood pressure	Smoking status	HbA1c	Urinary albumin	Serum creatinine	Cholesterol	Eyes	Feet



Independent Diabetes Trust
PO Box 294 Northampton NN1 4XS

**For further information about
all our FREE leaflets contact us:**

HELPLINE: 01604 622837
Email: enquiries@iddtinternational.org
www.iddtinternational.org